

SUPPORTING INFORMATION

Crystal Structures, Metal-activation and DNA-binding Properties of Two-domain IdeR from *Mycobacterium tuberculosis*

Goragot Wisedchaisri^{1⊥}, C. James Chou^{2⊥§}, Meiting Wu¹, Claudia Roach¹, Adrian E.
Rice^{1‡}, Randall K. Holmes³, Craig Beeson⁴, and Wim G. J. Hol^{1,5*}

† This work was supported by NIH grants CA65656 to W.G.J.H. and grant AI14107 to R.K.H.

¹Department of Biochemistry and Biomolecular Structure Center, University of Washington, Seattle, WA 98195, USA

²Department of Chemistry, University of Washington, Seattle, WA 98195

³Department of Microbiology, University of Colorado School of Medicine, Aurora, CO 80045, USA

⁴Department of Pharmaceutical Sciences, Medical University of South Carolina, Charleston, SC 29425, USA

⁵Howard Hughes Medical Institute, University of Washington, Seattle, WA 98195, USA

§Present address: Department of Molecular Biology, Scripps Research Institute, 10550 North Torrey Pines Road, MB-27, La Jolla, CA 92037, USA

‡Present address: Department of Biochemistry and Molecular Biophysics, California Institute of Technology, M/C 114-96, Pasadena, CA 91125, USA

⊥ G.W. and C.J.C. contributed equally to the work.

* To whom correspondence should be addressed. wghol@u.washington.edu
Tel: (206) 685-7044, Fax: (206) 685-7002

Table S1 Equilibrium constants for metal ion activation of 2D-IdeR for binding to the *fxbA* operator

Metal ion	$\langle K_d \rangle^a$ (μM)	$\langle K_d \rangle^a$ for full-length IdeR ^b (μM)
Co ²⁺	24.7 \pm 1.1	22.6 \pm 1.9
Ni ²⁺	4.3 \pm 1.0	13.1 \pm 0.1

^a The average value of at least three titrations.

^b Data from Chou *et al.* (14)

Table S2 Metal-ligand distances (Å) for all metal-binding sites

Site	Residue	Atom	2D-IdeR		2D-IdeR:DNA complex (crystal form I)				2D-IdeR:DNA complex (crystal form II)			
			A	B	A	B	C	D	A	B	C	D
Ni 1	His79	N ^{ε2}	2.22	2.08	2.04	2.03	2.04	2.03	2.02	2.01	2.01	2.02
	Glu83	O ^{ε2}	2.14	2.11	1.96	1.96	1.96	1.96	1.95	1.95	1.94	1.94
	His98	N ^{δ1}	2.15	2.11	2.04	2.05	2.04	2.03	2.02	2.02	2.03	2.00
	Phosphate	O ¹	2.17	2.25	-	-	-	-	-	-	-	-
	H ₂ O (1)	O	2.19	2.08	2.60	2.61	-	2.62	-	-	-	-
	H ₂ O (2)	O	2.37	2.19	-	-	-	-	-	-	-	-
	Acetate (1)	O	-	-	-	-	-	-	2.12	2.49	2.17	2.48
	Acetate (1)	O ^{XT}	-	-	-	-	-	-	2.35	2.45	2.27	1.87
Ni 2	Met10	S ^δ	2.42	2.47	2.37	2.33	2.36	2.34	2.34	2.33	2.35	2.34
	Cys102	O	2.15	2.08	1.97	1.96	1.95	1.95	1.96	1.95	1.94	1.95
	Cys102	S ^γ	-	-	2.79	2.61	2.85	3.08	2.75	2.73	2.78	2.81
	Cys102 ^a	O ¹	1.91	2.11	-	-	-	-	-	-	-	-
	Glu105	O ^{ε2}	2.04	2.12	1.95	1.95	1.96	1.95	1.94	1.94	1.94	1.94
	His106	N ^{ε2}	2.07	1.98	2.03	2.04	2.02	2.04	2.02	2.03	2.02	2.01
	H ₂ O	O	2.21	2.18	1.99	2.98	2.31	2.27	-	-	2.33	-
Ni 3	His61	N ^{δ1}	-	-	2.03	2.03	2.02	2.03	2.02	2.03	2.03	2.01
	H ₂ O	O	-	-	-	1.83	2.07	2.67	-	-	-	-
	H ₂ O	O	-	-	-	2.32	2.44	2.74	-	-	-	-
	H ₂ O	O	-	-	-	2.33	2.85	-	-	-	-	-
	Acetate (2)	O	-	-	-	-	-	-	2.21	1.93	2.39	2.42
	Acetate (2)	O ^{XT}	-	-	-	-	-	-	2.87	2.07	2.48	1.97
	Acetate (3)	O	-	-	-	-	-	-	2.08	-	2.66	-
	Acetate (3)	O ^{XT}	-	-	-	-	-	-	2.74	-	2.50	-
Na	Phosphate	O ^{2P}	-	-	2.86 ^b (F19)	-	3.24 ^b (E16)	3.17 ^b (F24)	-	-	-	-
	H ₂ O (1)	O	-	-	1.86	-	2.28	1.94	-	-	-	-
	H ₂ O (2)	O	-	-	2.00	-	2.49	2.12	-	-	-	-
	H ₂ O (3)	O	-	-	2.10	-	2.54	2.35	-	-	-	-
	H ₂ O (4)	O	-	-	2.24	-	-	-	-	-	-	-

^aCys 102 is modeled as S-hydroxy cystein in 2D-IdeR structure.^bPhosphate oxygen of residues in parenthesis coordinate to sodium ions.

Table S3 Distance (Å) between Ser37/Pro39 and interacting nucleotide bases (with a 3.9 Å cutoff distance) in the 2D-IdeR:DNA complex structures

IdeR subunit	A			B				C			D			
Nucleotide (chain)	T20 (F)	G21 (F)	C12 (E)	T22 (E)	A23 (E)	T10 (F)	T9 (F)	T17 (E)	G18 (E)	C15 (F)	T25 (F)	A26 (F)	T7 (E)	T6 (E)
Crystal form I														
Ser37 C ^β	3.49 (C5M)	-	-	3.55 (C5M)	-	-	-	3.42 (C5M)	-	-	3.47 (C5M)	-	-	-
Ser37 O ^γ	3.38 (C5M)	-	-	3.40 (C5M)	-	-	-	3.44 (C5M)	-	-	3.34 (C5M)	-	-	-
Pro39 N	-	-	-	-	-	-	3.50 (C5M)	-	-	-	-	-	-	3.74 (C5M)
Pro39 C ^α	-	-	-	-	-	-	3.80 (C5M)	-	-	-	-	-	-	-
Pro39 C ^β	3.09 (O4)	-	-	3.49 (O4)	-	3.46 (O4)	-	3.27 (O4)	-	-	3.41 (O4)	-	-	-
Pro39 C ^γ	2.97 (O4)	3.58 (O6)	3.68 (N4)	3.34 (O4)	3.51 (N6)	3.19 (O4)	3.70 (O4)	2.98 (O4)	3.87 (O6)	3.70 (N4)	3.26 (O4)	3.57 (N6)	3.47 (O4)	3.78 (O4)
Pro39 C ^δ	3.65 (C5M)	-	-	-	-	-	3.85 (O4)	3.71 (C5M)	-	-	-	-	-	-
	3.82 (C5M)	-	-	-	-	-	3.86 (C5M)	3.75 (C5M)	-	-	-	-	-	-
Crystal form II														
Ser37 C ^β	3.49 (C5M)	-	-	3.28 (C5M)	-	-	-	3.34 (C5M)	-	-	3.56 (C5M)	-	-	-
Ser37 O ^γ	3.57 (C5M)	-	-	3.35 (C5M)	-	-	-	3.29 (C5M)	-	-	3.22 (C5M)	-	-	-
Pro36 N	-	-	-	-	-	-	3.82 (C5M)	-	-	-	-	-	-	-
Pro39 C ^β	3.30 (O4)	-	3.85 (N4)	3.39 (O4)	-	3.49 (O4)	-	3.13 (O4)	-	-	3.37 (O4)	-	3.74 (O4)	-
Pro39 C ^γ	3.30 (O4)	3.89 (O6)	3.54 (N4)	3.33 (O4)	3.18 (N6)	3.40 (O4)	3.46 (O4)	3.18 (O4)	3.74 (O6)	3.71 (N4)	3.31 (O4)	3.42 (N6)	3.57 (O4)	-
	3.89 (C5M)	-	-	-	-	-	-	-	-	-	3.74 (C5M)	-	-	-
Pro39 C ^δ	-	-	-	-	-	-	3.76 (O4)	-	-	-	-	-	-	-

Table S4 Distance (Å) between selected atoms of Gln43 and interacting nucleotide bases
(with a 3.9 Å cutoff distance) in the 2D-IdeR:DNA complex structures

IdeR subunit	A		B		C^a		D	
Nucleotide (chain)	<i>C17</i> (F)	<i>C18</i> (F)	<i>C19</i> (E)	<i>C20</i> (E)	<i>G14</i> (E)	<i>G15</i> (E)	<i>T22</i> (F)	<i>G23</i> (F)
Crystal form I								
Gln43 C ^δ				3.70 (N4)				
Gln43 O ^{ε1}	-	3.83 (C4) 3.07 (N4) 3.71 (C5)	3.68 (C5)	3.50 (C4) 2.70 (N4) 3.48 (C5)	-	-	-	-
Gln43 N ^{ε2}	-	-	-	-	-	-	-	-
Crystal form II								
Gln43 C ^δ				3.86 (N4)				
Gln43 O ^{ε1}	3.65 (N7)	3.38 (N4) 3.81 (C5)	3.52 (C5)	3.66 (C4) 2.81 (N4) 3.65 (C5)	-	-	-	-
Gln43 N ^{ε2}	-	-	-	-	-	-	-	-

^aGln43 of subunit C in crystal form II makes the following unusual contacts to nucleotide bases:

Gln43 O^{ε1} - C16E N4 (3.23 Å) and Gln43 N^{ε2} - G17F O6 (3.35 Å)

Figure S1 Representative dynamic-light scattering data

